

# Nutrient & Manure Management Tables

**Table 1. Annual Manure Production and Nutrient Excretion from Livestock**

Animal Type	Manure Production per 1,000 lb. of Animal Weight		Excreted Nutrients in Manure per 1,000 lb. of Animal Weight		
	Solid tons/year	Liquid gal./year	N lb./year	P2O5 lb./year	K2O lb./year
<b>BEEF CATTLE</b>					
Calf	19.5	4,591	162	73	130
Finishing	9.0	2,141	131	39	83
Cow	16.8	3,982	128	66	106
<b>DAIRY CATTLE</b>					
Calf	14.6	3,358	146	24	122
Heifer	11.0	2,536	112	39	112
Lactating	20.3	4,876	263	135	146
Dry Cow	9.3	2,241	110	40	88
Veal	4.8	1,153	44	29	73
<b>SWINE</b>					
Nursery	13.9	3,358	292	146	146
Finishing	9.0	2,166	219	73	97
Gestating Sow	4.1	998	61	37	49
Lactating Sow	8.5	2,025	165	107	127
Boar	3.8	900	49	37	37
<b>POULTRY</b>					
Layer	9.1	2,068	316	97	146
Broiler	17.3	4,198	383	256	183
Turkey (F)	8.6	2,044	285	186	124
Turkey (M)	6.8	1,606	203	135	88
Duck	20.1	4,836	392	310	237
<b>HORSE</b>					
Pleasure	9.9	2,394	66	22	22
"Racer"	10.1	2,446	110	55	84
<b>SHEEP</b>					
Feeder	7.5	1,825	146	73	146

Adapted From: Manure Characteristics, MWPS-18 Section 1, Midwest Plan Service, 2004 Second Edition

**Table 2. Nitrogen Losses From Animal Manure as Affected by Method of Handling and Storage**

Manure Storage and Handling method	Manure Type	% Storage Nitrogen Loss
Daily scrape and haul	Solid (tons)	25
Manure pack	Solid (tons)	30
Open lot	Solid (tons)	50
Litter	Solid (tons)	35
Above ground tank	Liquid (gallons)	20
Below ground covered pit	Liquid (gallons)	20
Below ground open pit	Liquid (gallons)	25
Under-floor dry storage	Solid (tons)	25
Under-floor liquid storage	Liquid (gallons)	20
Earthen storage	Liquid (gallons)	30
Lagoon	Liquid (gallons)	75

Adapted From: Animal Manure as a Plant Nutrient Source, ID-101, Cooperative Extension Service, Purdue University, 2001

**Table 3. Nutrient Content of Stored Manure**

Animal Type	Liquid Manure (lb./1000 gallons)			Solid Manure (lb./Tons)		
	N	P2O5	K2O	N	P2O5	K2O
<b>BEEF</b>						
Cows	20	16	24	7	4	7
Finishing Cattle	29	18	26	11	7	11
<b>DAIRY</b>						
Cows	31	15	19	10	3	6
Heifers	32	14	28	10	3	7
<b>SWINE</b>						
Farrowing	15	12	11	14	6	4
Nursery	25	19	22	13	8	4
Gestation	25	25	24	9	7	5
Finishing	58	44	40	16	9	5
<b>POULTRY</b>						
Broilers	63	40	29	46	53	36
Layers	57	52	33	34	51	26
Tom Turkeys	53	40	29	40	50	30
Hen Turkeys	60	38	32	40	50	30
<b>HORSE</b>						
				14	4	14
<b>SHEEP</b>						
				18	11	26

Manure Management in Minnesota, FO-3553-C, University of Minnesota Extension, 2012 Manure Characteristics, MWPS-18 Section 1, Midwest Plan Service, 2004

**Table 4. Nitrogen Availability and Loss as Affected by Method of Manure Application and Animal Species**

Year Available (1)	% of Total Nitrogen Available Per Year				
	Broadcast Incorporation Time (2)			Injection	
	> 96 hrs	12-96 hrs.	<12 hrs.	Sweep	Knife
<b>BEEF</b>					
Year 1	25	45	60	60	50
Year 2	25	25	25	25	25
Lost	40	20	5	5	10
<b>DAIRY</b>					
Year 1	20	40	55	55	50
Year 2	25	25	25	25	25
Lost	40	20	10	5	10
<b>SWINE</b>					
Year 1	35	55	75	80	70
Year 2	15	15	15	15	15
Lost	50	30	10	5	15
<b>POULTRY</b>					
Year 1	45	55	70	NA	NA
Year 2	25	25	25	NA	NA
Lost	30	20	5		

From: Manure Management in Minnesota, FO-3553-C, University of Minnesota Extension Service, 2007

(1) Third year available N is not listed but can be computed by adding 1st and 2nd year and lost percentages and subtracting this sum from 100.

(2) Timing categories refer to the length of time between manure application and incorporation.

## Calibrating Your Manure Spreader

- Determine manure weight (solid manure) or manure volume (liquid manure) per spreader load (Use measured manure weight or 90% of the manufacturer's listed volume for liquid).
- Calculate rate based on loads applied per field of known size **OR** calculate rate based on acres covered per load ( $Length \times Width \text{ of Spread (ft.}^2\text{)}/43,560$ ).



**Table 5. Average Nutrient Removal Rates for Crops in the Northcentral Region**

Crop	Crop Nutrient Removal (lb. per unit)			
	Yield Units	N	P2O5	K2O
Alfalfa	ton (air dry)	51	12	49
Alsike clover	ton (air dry)	41	11	54
Barley (grain)	bushel	0.99	0.4	0.32
Barley (grain & straw)	bushel	1.39	0.56	1.52
Birdsfoot trefoil	ton (air dry)	45	11	42
Canola	bushel	1.9	1.2	2
Corn (grain)	bushel	0.9	0.38	0.27
Corn silage	ton (as fed)	9.7	3.1	7.3
Beans, dry	bushel	3	0.79	0.92
Bromegrass	ton	32	10	46
Orchardgrass	ton	36	13	54
Oats (grain)	bushel	0.77	0.28	0.19
Oats (straw)	ton	12	6.3	37
Potatoes (tuber)	cwt	0.32	0.12	0.55
Red clover	ton	45	12	42
Rye (grain)	bushel	1.4	0.46	0.31
Rye (straw)	ton	12	3	22
Soybeans	bushel	3.8	0.84	1.3
Sugar beets	ton	3.7	2.2	7.3
Sunflowers	cwt	2.7	0.97	0.9
Wheat (grain)	bushel	1.5	0.6	0.34
Wheat (straw)	ton	14	3.3	24

Source: International Plant Nutrition Institute (IPNI) Sept. 2005 <http://nanc.ipni.net/articles/NANCO005-EN>

**Table 8. Legume Nitrogen Credits**

For Corn, Wheat, and Barley grown the 1st and 2nd year after a legume crop

Legume (Previous Crop)	Corn		Wheat & Barley	
	1st year	2nd year	1st year	2nd year
Soybeans	40	0	20	0
Edible beans	20	0	10	0
Field peas	20	0	10	0
Red Clover	75	35	35	20
Harvested sweet clover	20	0	10	0
Harvested alfalfa or nonharvested sweet clover (plants/ft 2)				
4 or more	150	75	75	35
2-3	100	50	50	25
1 or less	40	0	0	0

**Table 6. Common Fertilizer Analyses**

Example to calculate fertilizer price per pound:  
 Urea (46-0-0) = \$600/ton  
 (2000 lb. X .46% N) = 920 lb. N/ton  
 \$600 / 920 lb. = \$0.65/lb.

Fertilizer	Analysis
<b>N</b>	
Anhydrous Ammonia	82-0-0
Ammonium Nitrate	34-0-0
Urea	46-0-0
UAN Solution (Urea Ammonium Nitrate)	28 to 32-0-0
Ammonium Sulfate	21-0-0-24(S)
<b>P</b>	
Triple Superphosphate (TSP)	0-44 to 0-46
Diammonium Phosphate (DAP)	18-46-0
Monoammonium Phosphate (MAP)	11-52-0
Ammonium Polyphosphate Liquid (APP)	10-34-0
Ammonium Polyphosphate Dry (APP)	15-62-0
<b>K</b>	
Potassium Chloride (Muriate of Potash)	0-0-60
Potassium Sulfate	0-0-50-18(S)
Potassium-Magnesium Sulfate (Sul-fo-mag)	0-0-22-22(S)-11(Mg)
Potassium Nitrate	13-0-44

**Table 7. Nitrogen Rate Guidelines for Corn (When Using a Manure Source)**

Previous Crop <sup>1</sup>	Soil / Field Productivity Potential	
	Highly Productive Soils	Medium Productive Soils <sup>2</sup>
	----- lb. N/A -----	
<b>CORN</b>	<b>130 - 180</b>	<b>130</b>
<b>SOYBEAN</b>	<b>100 - 140</b>	<b>100</b>

<sup>1</sup>For previous crops other than corn or soybeans use the corn following corn rate guideline and subtract any previous crop N credits in Table 8.

<sup>2</sup>Soil and environmental conditions that limit crop production such as erosion, poor soil drainage, restriction to root growth, short growing season, and marginal growing season rainfall, among others, would qualify a site as having medium productivity potential.

**Table 9. Nitrogen Sources Per Pound Conversions**

Anhydrous Ammonia (82-0-0)		Urea (46-0-0)		UAN (28-0-0)	
Price per ton	N Price per lb	Price per ton	N Price per lb	Price per ton	N Price per lb
400	\$0.25	350	\$0.38	200	\$0.36
450	\$0.28	385	\$0.42	225	\$0.40
500	\$0.30	420	\$0.46	250	\$0.45
550	\$0.34	455	\$0.49	275	\$0.49
600	\$0.37	490	\$0.53	300	\$0.54
650	\$0.40	525	\$0.57	325	\$0.58
700	\$0.43	560	\$0.61	350	\$0.63
750	\$0.46	595	\$0.65	375	\$0.67
800	\$0.49	630	\$0.68	400	\$0.71
850	\$0.52	665	\$0.72	425	\$0.76
900	\$0.55	700	\$0.76	450	\$0.80
950	\$0.58	735	\$0.80	475	\$0.85
1,000	\$0.61	770	\$0.84	500	\$0.89
1,050	\$0.64	805	\$0.88	525	\$0.94
1,100	\$0.67	840	\$0.91	550	\$0.98
1,150	\$0.70	875	\$0.95	600	\$1.07
1,200	\$0.73	910	\$0.99	650	\$1.16

**Table 10. Conversion Factors**

1 acre = 43,560 sq. ft	
1 cubic ft. = 7.48 gallons	
1 gallon of water = 8.33 lb.	
1 ton = 2000 lb.	
<b>SOIL TESTING CONVERSIONS</b>	
Plow Layer (6-7 in) = ppm X 2 = lb./acre	
Top 12 inches = ppm X 4 = lb./acre	
Top 24 inches = ppm X 8 = lb./acre	
P2O5 X 0.44 = P	
P X 2.29 = P2O5	
K2O X 0.83 = K	
K X 1.20 = K2O	
<b>FERTILIZER CONVERSIONS</b>	
1 gal. of UAN (28%) = 10.66 lb.	
1 gal. (10-34-0) = 11.65 lb.	
1 gal. (7-21-7) = 11.0 lb.	
1 gal. (9-18-9) = 11.11 lb.	

**Manure Management Related Websites**

- Certified Manure Testing Laboratories: <http://www2.mda.state.mn.us/webapp/lis/manurelabs.jsp>
- University of Minnesota Manure Management: <http://www.manure.umn.edu/>
- Minnesota Department of Ag Manure Management: <http://www.mda.state.mn.us/protecting/conservation/practices/manuregmt.aspx>
- Minnesota NRCS Manure Management: <http://www.mn.nrcs.usda.gov/technical/ecs/nutrient/manure/manure.htm>
- Nitrogen Best Management Practices (BMPs): <http://www.mda.state.mn.us/nitrogenbmps>
- Nitrogen Rate Calculator: <http://extension.agron.iastate.edu/soilfertility/nrate.aspx>
- Minnesota Pollution Control Agency: <http://www.pca.state.mn.us/index.php/topics/feedlots/feedlot-nutrient-and-manure-management.html?menuid=&redirect=1>

In accordance with the Americans with Disabilities Act, this information is available in alternative forms of communication upon request by calling 651/201-6000. TTY users can call the Minnesota Relay Service at 711 or 1-800-627-3529. The MDA is an equal opportunity employer and provider.

Printing made possible through a grant from Conservation Marketplace Midwest